

Indicators	Method	Primary target group/How to access/Who collect the data	How often	Who compiled and analyzes	Who receive and act on the information
Prevalence of whole households that purchase/use fortified flour (or bread)	Subjects are asked about the type of flour they purchase using a standard questionnaire	Non pregnant Women in childbearing age, interview in sentinel health facility (SAAJ e PF), health staff trained	First round, and every 6 months	Nutrition Department/MoH	CONFAM and sentinel facilities
Per capita consumption of fortified flour	Subjects report household size and quantity of flour/flour products purchased in specified period	NPWCBA, interview in household, SETSAN Staff trained	First round, every year	SETSAN	CONFAM and SETSAN
Prevalence of Anemia	Blood Hb test	NPWCBA, health center (SAAJ e PF), health staff trained	First month of start of FF, and next every year	Dept. Health Information/MoH	CONFAM and sentinel facilities
Prevalence of NTD	Health facilities report NTD births	All babies born in Provincial Hospitals, maternity staff	First month of start of FF, and next every year	Dept. Health Information/MoH	CONFAM and Provincial hospitals
✓ Prevalence of Anemia	Blood tests	Non pregnant Women in childbearing age and children under 5 years old	First round before start the FF Program and ~1year before achieve 80% of household consuming fortified flour	M&S team of CONFAM	CONFAM, government and other stakeholders at all levels
✓ Prevalence of Fe deficiency (TFR, Ferritin)					
✓ Inflammatory indicators					
✓ Prevalence of B6, B12					
✓ Prevalence of folate sufficiency					
✓ Percentage of population	Subjects report household size and	Non pregnant Women in childbearing age and		M&S team of CONFAM	

consuming fortified flours	quantity of flour/flour products purchased/used	children under 5 years old			
-------------------------------	---	----------------------------	--	--	--